

BRL R 1463

BRL

AD-1000000

REPORT NO. 1463

NEGATIVE ION REACTIONS IN NO-H₂O MIXTURES

by

L. J. Puckett
W. C. Lineberger

December 1969

This document has been approved for public release and sale;
its distribution is unlimited.

U.S. ARMY ABERDEEN RESEARCH AND DEVELOPMENT CENTER
BALLISTIC RESEARCH LABORATORIES
ABERDEEN PROVING GROUND, MARYLAND

Destroy this report when it is no longer needed.
Do not return it to the originator.

The findings in this report are not to be construed as
an official Department of the Army position, unless
so designated by other authorized documents.

B A L L I S T I C R E S E A R C H L A B O R A T O R I E S

REPORT NO. 1463

DECEMBER 1969

NEGATIVE ION REACTIONS IN NO-H₂O MIXTURES

L. J. Puckett

W. C. Lineberger

Signature and Propagation Laboratory

This document has been approved for public release and sale;
its distribution is unlimited.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183

BALLISTIC RESEARCH LABORATORIES



Report No. 1463

LJPuckett/WCLineberger†/n11
Aberdeen Proving Ground, Md.
December 1969

NEGATIVE ION REACTIONS IN NO-H₂O MIXTURES*

ABSTRACT

A stationary afterglow system has been utilized to determine rate constants for thermal energy negative ion-molecule reactions in photo-ionized NO-H₂O mixtures. When the decay of the plasma is controlled by ambipolar diffusion of positive and negative ions quantitative determination of rate constants is shown to be feasible. The plasma transition from electron-positive ion ambipolar diffusive domination of the transport loss processes to domination by positive ion-negative ion ambipolar diffusion is explained by a model which includes the effects of negative ion trapping. Prominant negative ions in the afterglow include NO₂⁻, its hydrates, and clusters involving HNO₂. Reaction rate constants for the processes



are found to be $1.3 \pm 0.3 \times 10^{-28} \text{ cm}^6/\text{sec}$ and $3.4 \pm 1.3 \times 10^{-29} \text{ cm}^6/\text{sec}$ at 293 K, respectively. Steady glows in NO-H₂O-O₂ mixtures revealed that NO₃⁻ and the impurity HCO₃⁻ also formed multiple hydrates and clustered with HNO₂. These results indicate that the terminal negative ions in the D-region of

the ionosphere will likely be hydrated.

* Research supported in part by Defense Atomic Support Agency

† Present Address: Joint Institute for Laboratory Astrophysics
University of Colorado
Boulder, Colorado 80302

TABLE OF CONTENTS

	Page
ABSTRACT	3
LIST OF ILLUSTRATIONS	7
I. INTRODUCTION	9
II. EXPERIMENTAL APPARATUS AND PROCEDURES	10
III. AFTERGLOW ANALYSIS	12
IV. EXPERIMENTAL RESULTS ANALYSIS AND DISCUSSION	17
DISTRIBUTION LIST	27

LIST OF ILLUSTRATIONS

Figure		Page
1.	Schematic diagram of stationary afterglow apparatus . . .	11
2.	Temporal afterglow profiles of NO^+ , $\text{NO}^+\cdot\text{NO}$ and NO_2^- wall currents at 50 mTorr NO pressure	13
3.	Evolution of the negative ion spectrum as a function of increasing H_2O concentration in NO at 400 mTorr NO pressure. The integrated spectrum for each H_2O concentration is normalized to the same value .2.	18
4.	Variation of NO_2^- reactive loss frequency as a function of the product of NO and H_2O pressures	22
5.	Negative ion spectrum in 200 mTorr NO, 5 mTorr H_2O and 5 mTorr O_2	23

I. INTRODUCTION

Recently Lineberger and Puckett^{1,2*} reported stationary afterglow measurements of NO^+ reactions leading to the formation of $\text{NO}^+\cdot\text{NO}$, $\text{NO}^+\cdot n(\text{H}_2\text{O})$ and $\text{H}_3\text{O}^+\cdot n(\text{H}_2\text{O})$ ions in photoionized $\text{NO-H}_2\text{O}$ mixtures. These investigations elucidated a mechanism by which NO^+ ions can be lost in reactions with atmospheric water vapor. As a consequence of these reactions it is understandable that NO^+ should not be regarded as a terminal positive ion in the D-region of the ionosphere.

Ferguson³ and LeLevier and Branscomb⁴ have reviewed D-region negative ion chemistry and **concluded that the "terminal" negative ions are NO_2^- and NO_3^-** . This conclusion was based on the observation that these ions are formed through chain-breaking reactions that do not permit the electron to be freed again. Therefore, in their context, "terminal" implies that the ions are indestructable except through ion-ion mutual neutralization processes. We report results which demonstrate that both NO_2^- and NO_3^- ions do, however, undergo clustering reactions with H_2O and HNO_2 at 293 K.

The negative ion-molecule reaction rate constants reported in this paper are the first such measurements known to the authors to be made using stationary afterglow techniques. In order to obtain quantitative negative ion reaction rate information from a stationary afterglow, it is necessary to make observations subsequent to the disappearance of electrons from the decaying plasma during the interval when positive

*References are listed on page 25.

ion-negative ion ambipolar diffusion is the dominant transport loss mechanism. The transition from positive ion-electron ambipolar diffusive domination to positive ion-negative ion ambipolar diffusive domination is marked by a sudden increase in the negative ion wall current, and a sudden decrease in the positive ion wall current. A model is presented which accounts for the features of this transition.

II. EXPERIMENTAL APPARATUS AND PROCEDURES

The basic apparatus employed in this experiment is the photoionized-stationary-afterglow instrument described previously¹ and only a brief account of the apparatus will be presented here. For the present work the mass filter was modified to permit observation of negative ions. A schematic diagram of the apparatus is shown in Fig. 1.

The afterglow cavity is an ultra-high-vacuum, bakeable, gold-coated-stainless-steel cylinder 18 inches in diameter and 36 inches long. Information on the individual ion species in the plasma afterglow is obtained by means of time-resolved mass spectrometry of the ions which pass through a 0.60 mm diameter sampling orifice in the cavity wall. The sampling orifice is contained in a plate which is contoured to the shape of the cavity wall and electrically insulated from the wall. The potential on the plate is set at a variable but low attractive potential ($<100\text{mV}$). The orifice plate potential did not effect the rate constant determinations in this work; however, we have noted¹ that observed diffusion loss rates are affected by the plate potential. Consequently, care must be exercised in all measurements of diffusion coefficients in cases where draw-out




Figure 1. Schematic diagram of stationary afterglow apparatus

potentials are employed.

In this investigation the negative ions were formed by electron attachment in the gas. The electrons were produced by photoionization of NO by means of krypton resonance radiation (123.6 and 116.5 nm) from a pulsed microwave-powered discharge lamp. Initial ion density is sufficiently low ($\sim 10^6 \text{ cm}^{-3}$) that recombination loss rates are negligible compared with reactive and diffusive loss rates.

The NO gas used in this work was processed in the following manner. Specially prepared gas of 99.9 per cent stated purity was obtained from a steel cylinder. The gas was passed through a stainless steel and glass line to a LN_2 trap where it was condensed. By means of a refrigerating vapor bath⁵ the trap temperature could be maintained within ± 1.0 K of any desired temperature in the range 77 to ~ 300 K. The NO vapor at the selected trap temperature was passed through a servo-controlled leak valve to the afterglow chamber. The experimental results in this paper were found to be insensitive to trap temperatures below ~ 200 K. Above this temperature trace amounts of NO_2 in the NO were not completely trapped and appeared as impurity ions in the afterglow. Water vapor densities required for the rate constant determinations were obtained in the manner described previously².

III. AFTERGLOW ANALYSIS

Fig. 2 shows the temporal profiles of the principal positive and negative ions in a photoionized NO afterglow at a total pressure of 50 mTorr. The primary ion NO^+ and its termolecular reaction to form $\text{NO}^+ \cdot \text{NO}$




Figure 2. Temporal afterglow profiles of NO^+ , $\text{NO}^+ \cdot \text{NO}$ and NO_2^- wall currents at 50 mTorr NO pressure

have been discussed previously¹. The principal negative ion observed was NO_2^- . The initial formation mechanism for NO_2^- is currently under investigation and will be reported in a future publication.

The temporal profiles in Fig. 2 show marked transitions in positive and negative ion behavior at ~ 170 msec. The transitions in the ion currents to the wall are associated with the transition in volume transport from electron-ion ambipolar diffusive domination to positive ion-negative ion ambipolar diffusive domination, and the resulting release of trapped negative ions. The trapping of negative ions prior to this transition, coupled with sampling discrimination effects, have thwarted previous efforts⁶ to measure negative ion-molecule reaction rates in stationary afterglow experiments. In order to establish the validity of the reaction rate data presented here, it is necessary to investigate and explain the features of the transition. This section is accordingly devoted to an analysis of the plasma decay for times before, during, and after the transition.

In Fig. 2, for times < 170 msec, the positive ion decays are completely explained¹ by interconversion and positive ion-electron ambipolar diffusive⁷ loss. The principal features of the plasma decay that remain to be explained are as follows:

1. The negative ion wall current increases sharply at 170 msec and then decays exponentially.
2. All ions decay exponentially with the same time constant after 170 msec.
3. Both of the positive ion wall currents exhibit a sudden decrease

by approximately a factor of two at 170 msec.

These features are all quantitatively explained in the context of a simple ambipolar diffusion model, as outlined below. The diffusion current density, Γ , of positive ions, negative ions and electrons in the afterglow may be expressed by⁷

$$\Gamma_+ = -D_+ \nabla N_+ + N_+ \mu_+ E \quad (1)$$

$$\Gamma_- = -D_- \nabla N_- - N_- \mu_- E \quad (2)$$

and

$$\Gamma_e = -D_e \nabla N_e - N_e \mu_e E \quad (3)$$

respectively. The quantities D_i and μ_i are free diffusion coefficients and mobilities, while E is the electric field produced by non-charge neutrality in the plasma. The free diffusion term, $-D_i \nabla N_i$, is due to the density gradient of the i th charged species, while the mobility term, $N_i \mu_i E$, describes the field induced charged particle drift in the gas. Although there are no applied electric fields in the afterglow, a "self-field" develops due to the initial rapid diffusion of electrons, producing a net charge separation which retards electron diffusion and enhances positive ion diffusion. It may be shown⁷ using the Einstein relation, $\mu = \frac{eD}{kT}$, that the free diffusion terms and the mobility terms are of equal magnitudes. For positive ions the mobility term produces a current in the same direction as that of free diffusion.

Hence,
$$\Gamma_+ = -2D_{+,e} \nabla N_+ \equiv -D_{+,e} \nabla N_+, \quad (4)$$

where $D_{+,e}$ is defined as the positive ion-electron ambipolar diffusion coefficient, and is equal to twice the free diffusion coefficient, D_+ .

In the case of electrons, the free diffusion coefficient, D_e , and the mobility, μ_e , are a factor of 10^5 greater than those corresponding

to ions in the afterglow. In spite of the fact that diffusion and mobility terms are in opposition for electrons, these terms are of sufficient magnitude such that a departure of about 1 part in 10^5 from complete cancellation is adequate to maintain $\Gamma_e = \Gamma_+$.

In addition to being lost through positive ion-electron diffusion, electrons are lost in attaching reactions which form negative ions. The electric field opposes the radial diffusion of negative ions, however, and as a result

$$\Gamma_- \approx 0, \quad (5)$$

as can be seen in Fig. 2 for $t < 170$ msec. In this sense the negative ions are trapped in the afterglow. Thus, the plasma decays through positive ion-electron ambipolar diffusion until the number density of electrons is no longer sufficient to maintain the electric field which gave rise to ambipolar diffusion and negative ion containment. The collapse of the electric field in the plasma is evidenced in the afterglow profile at ≈ 170 msec. According to Eq.(3), in the absence of the electric field, $\Gamma_e = -D_e \nabla N_e$ and the remaining electrons are lost very rapidly by free diffusive processes. In this simple model the positive ion diffusion current Γ_+ decreases from $2D_+ \nabla N_+$ to $D_+ \nabla N_+$, which gives rise to a rapid drop of wall current by a factor of two followed by a continued reduction by a factor of two in the exponential rate of plasma decay. Both of these features are apparent in the afterglow profile. The collapse of the E-field also terminates the containment of the negative ions in the afterglow and produces a rapid increase of Γ_- from ≈ 0 to the value $\Gamma_- = -D \nabla N_-$, which in the first approximation is equal Γ_+ . These characteristics in the negative ion behavior are also observable in the afterglow profile.

The relatively simple model described above qualitatively accounts for all of the observed transition features. A more refined numerical analysis of the transition, involving only the assumption of charge neutrality, has recently been completed by Kregel⁸. This more refined calculation reproduces both the observed buildup of negative ion current prior to the transition, and the smoothing of the transition observed in the positive ion wall current.

IV. EXPERIMENTAL RESULTS AND DISCUSSION

These measurements are believed to constitute the first measurements of negative ion reaction rate constants to be obtained with a stationary afterglow apparatus. Negative ions have been observed in other stationary afterglow experiments⁶, but because of small signals and ion discrimination effects, the previous observations were unable to follow the afterglow into the positive ion-negative ion ambipolar diffusion regime. Consequently for all times during the observations⁶ there were both negative ion sources which were difficult to evaluate and non-equilibrium ionic spatial distributions. As a result of these conditions meaningful negative ion reaction rate constants could not be obtained.

With the apparatus employed in this work the afterglow has been observed over seven decades of decay, four decades of which followed the electron-ion to ion-ion transition, i.e. four decades of decay in which there were negligible net sources of negative ions, and the ionic spatial distribution was a fundamental mode diffusive distribution.

Fig. 3 shows the negative ion spectrum in pure nitric oxide and in






Figure 3. Evolution of the negative ion spectrum as a function of increasing H_2O concentration in NO at 400 mTorr NO pressure. The integrated spectrum for each H_2O concentration is normalized to the same value

nitric oxide with varying amounts of water vapor. The dominant ion is NO_2^- followed in intensity by Cl^- . (The Cl probably originates in the AgCl cement used to attach the MgF_2 windows to the afterglow chamber.) In pure NO, NO_2^- is the dominant negative ion throughout the afterglow; however, when H_2O is added in small amounts other ions become prominent. Principal among these ions is $\text{NO}_2^- \cdot \text{HNO}_2$. Clusters of NO_2^- with H_2O become increasingly important as the water vapor pressure increases. The hydrated NO_2^- and Cl^- are formed through the following reactions:



The rate constants k_1 and k_2 can be deduced through the following analysis.

If there are no NO_2^- sources and the dominant loss processes for NO_2^- ions in dilute H_2O -NO mixtures are positive ion-negative ion ambipolar diffusion and the reaction represented by Eq.(6), then for a fundamental mode diffusive distribution in a cylindrical cavity of radius R cm, the NO_2^- density in the afterglow may be expressed as

$$[\text{NO}_2^-(r,t)] = [\text{NO}_2^-(o,T)] J_0(2.405r/R) \cdot \exp\left\{-\left(\frac{D^+ + D^-}{\Lambda^2}\right) - k_1[\text{NO}][\text{H}_2\text{O}]\right\}(t-T), \quad (8)$$

which is valid for $t > T$. Brackets, $[\]$, denote the number density in cm^{-3} , $[\text{NO}_2^-(o,T)]$ is the axial number density at the time, T, of the transition from electron-ion to ion-ion ambipolar diffusion domination, and Λ is the characteristic diffusion length of the afterglow chamber.

It can be shown that under the proper experimental conditions¹, the count rate of a mass-analyzed ionic species is directly proportional to the ionic volume number density of that species. If the reciprocal time

constant for the observed decay of NO_2^- is denoted by ν , then

$$\nu = \frac{D_{+,-}}{\Lambda^2} + k_1 [\text{NO}] [\text{H}_2\text{O}] \quad (9)$$


When trace amounts of H_2O are added to the NO , $\frac{D_{+,-}}{\Lambda^2}$ is not significantly affected and the contribution of $k_1 [\text{NO}] [\text{H}_2\text{O}]$ to ν can be measured. A plot of $\nu - \frac{D_{+,-}}{\Lambda^2}$ as a function of $[\text{NO}] [\text{H}_2\text{O}]$ will indicate the dependence of the reaction on the NO and H_2O concentrations, and from this information the rate constant k_1 can be evaluated. The experimental data shown in Fig. 4 yield a value of $k_1 = 1.3 \pm 0.3 \times 10^{-28} \text{ cm}^6/\text{sec}$. Employing the same analysis for the Cl^- reactions as that described for NO_2^- the rate constant for the hydration of Cl^- , Eq. (7), was determined to be $k_2 = 3.4 \pm 1.3 \times 10^{-29} \text{ cm}^6/\text{sec}$. In order to deduce the sequence of reactions which produce the prominent ion, $\text{NO}_2^- \cdot \text{HNO}_2$, the following observations were made:

1. The NO_2^- count rate did not vary with either irradiation time (for times \gg the characteristic lifetime of the ion in the system) or with the residence time of gas in the chamber.
2. The count rate of $\text{NO}_2^- \cdot \text{HNO}_2$, however, did increase as a function of irradiation time, but did not increase with residence time of the gas in the chamber.


Observations 1 and 2, together, indicate that the reactant HNO_2 was produced through radiation chemistry. This conclusion is in accord with previous investigations² which delineated a source of HNO_2 through $\text{NO}^+ - \text{H}_2\text{O}$ gas phase chemistry.

3. The exponential decay of $\text{NO}_2^- \cdot \text{HNO}_2$ in the afterglow is substantially slower than that of NO_2^- , and lends further support to the conclusion that $\text{NO}_2^- \cdot \text{HNO}$ is not produced in reactions of NO_2^- with the


chamber walls, but rather in gas phase reactions. Fig. 4 of this paper reveals, however, that the dominant loss of NO_2^- in $\text{NO-H}_2\text{O}$ mixtures is the reaction



and not directly through



eventhough Fig. 3 indicates that the abundance of $\text{NO}_2^- \cdot \text{HNO}_2$ greatly exceeds that of $\text{NO}_2^- \cdot \text{H}_2\text{O}$. The indicated conclusion is therefore, that the principal source of $\text{NO}_2^- \cdot \text{HNO}_2$ is the reaction



with reaction (7) being a minor source of $\text{NO}_2^- \cdot \text{HNO}_2$ under the conditions of this investigation. Reaction (8) is an example of the "switching" reactions recently reported by Adams et al⁹, and the fact that reaction (8) is rapid indicates that the $\text{NO}_2^- \cdot \text{HNO}_2$ bond strength is greater than the $\text{NO}_2^- \cdot \text{H}_2\text{O}$ bond strength.

At the higher H_2O concentrations in Fig. 3 clustering reactions with both H_2O and HNO_2 prevail and the final negative ions are considerably more complex than NO_2^- . Similarly, clusters of the impurity, HCO_3^- , were observed to account for a large portion of the total ion spectrum at the higher water concentrations shown in Fig. 3. Further investigations revealed that NO_3^- present in $\text{NO-H}_2\text{O-O}_2$ mixtures (Fig. 5) also clustered with H_2O and HNO_2 . In this figure tentative identifications are made based on mass-to-charge ratios.

The present findings serve to indicate that those ions previously designated as terminal negative ions in the D-region of the ionosphere




Figure 4. Variation of NO_2^- reactive loss frequency as a function of the product of NO and H_2O pressures




Figure 5. Negative ion spectrum in 200 mTorr NO, 5 mTorr H₂O and 5 mTorr O₂

will certainly be hydrated, and perhaps clustering reactions with other D-region constituents will also be observed.

ACKNOWLEDGEMENTS

The authors have profited from many stimulating discussions of the plasma transition with Dr. M. D. Kregel. M. W. Teague assisted in acquisition and interpretation of much of the experimental data. Appreciation is due Dr. F. E. Niles and Dr. G. E. Keller for numerous discussions concerning experimental results.

REFERENCES

1. W. C. Lineberger and L. J. Puckett, "Positive Ions in Nitric Oxide Afterglows", Phys. Rev., to be published Oct. 1969.
2. W. C. Lineberger and L. J. Puckett, "Hydrated Positive Ions in Nitric Oxide-Water Afterglows", Phys. Rev., to be published.
3. E. E. Ferguson, Rev. Geophys., 5, 305 (1967).
4. R. E. Lelevier and L. M. Branscomb, J. Geophys. Res., 73, 27 (1968).
5. L. J. Puckett, M. W. Teague and D. G. McCoy, "Refrigerating Vapor Bath", Rev. Sci. Instr. submitted for publication.
6. W. L. Fite and J. A. Rutherford, Faraday Soc. Discussions, 37, 192 (1964).
7. H. J. Oskam, Phillips Res. Rpt., 13, 401 (1958).
8. M. D. Kregel, "Diffusion in Decaying Plasmas", J. Appl. Phys., to be published.
9. N. G. Adams, K. K. Bohme, D. B. Dunkin, F. C. Fehsenfeld, and E. E. Ferguson, "Flowing Afterglow Studies of Formation and Reactions of Cluster Ions of O_2^+ , O_2^- and O^- ", J. Chem. Phys., to be published.

DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
20	Commander Defense Documentation Center ATTN: TIPCR Cameron Station Alexandria, Virginia 22314	1	Commanding Officer U. S. Army Mobility Equipment Research & Development Center ATTN: Tech Docu Cen, Bldg 315 Fort Belvoir, Virginia 22060
1	Director Advanced Research Projects Agency ATTN: Dr. D. Mann Department of Defense Washington, D. C. 20301	1	Office of Vice Chief of Staff ATTN: CSAVCS-W-TIS Department of the Army Washington, D. C. 20310
1	Institute for Defense Analyses ATTN: Dr. E. Bauer 400 Army-Navy Drive Arlington, Virginia 22202	1	Commanding Officer U. S. Army Research Office (Durham) ATTN: Dr. Robert Mace Box CM, Duke Station Durham, North Carolina 27706
1	Director Defense Atomic Support Agency ATTN: STRA (RAEL) Dr. C. A. Blank Washington, D. C. 20305	3	Commander U. S. Naval Air Systems Command ATTN: AIR-604 Washington, D. C. 20360
1	DASA Information and Analysis Center TEMPO, General Electric Company ATTN: Dr. D. Griesinger 816 State Street Santa Barbara, California 93102	3	Commander U. S. Naval Ordnance Systems Command ATTN: ORD-9132 Washington, D. C. 20360
1	Commanding General U. S. Army Materiel Command ATTN: AMCRD-TE Washington, D. C. 20315	1	AFWL (WLREH, CPT Greene) Kirtland AFB New Mexico 87117
1	Commanding General U. S. Army Materiel Command ATTN: AMCRD-BN, Mr. N. Stulman Washington, D. C. 20315	6	AFCRL (CRUB, Dr. Keneshea, Dr. Hunt, Dr. Champion, Dr. Paulson, Dr. Murad; CRUS, Dr. Huffman) L. G. Hanscom Field Bedford, Massachusetts 01731
1	Commanding General U. S. Army Materiel Command ATTN: AMCRD-TP Washington, D. C. 20315	2	Director National Bureau of Standards ATTN: Dr. L. Kieffer Dr. S. Smith Washington, D. C. 20234

DISTRIBUTION LIST

<u>No. of</u> <u>Copies</u>	<u>Organization</u>	<u>No. of</u> <u>Copies</u>	<u>Organization</u>
8	Director Environmental Science Services Administration ATTN: Div 540.00, Dr. Fehsenfeld Dr. E. Ferguson Div 540.30, Dr. Albritton Dr. Schmeltekopf Div 224.00, Dr. G. Dunn Dr. W. C. Lineberger U. S. Department of Commerce Boulder, Colorado 80302	3	Lockheed Missiles and Space Company ATTN: Dr. R. Gunton Dr. R. Meyerott Dr. R. Varney 3251 Hanover Street Palo Alto, California 94304
1	AVCO-Everett Research Laboratory ATTN: Dr. E. Sutton 2385 Revere Beach Parkway Everett, Massachusetts 02149	1	Midwest Research Institute ATTN: Dr. T. A. Milne Kansas City, Missouri 64110
1	G. C. Dewey Corporation ATTN: Dr. M. Hirsh 202 East 44th Street New York, New York 10017	1	Pennsylvania State University Ionosphere Research Laboratory Electrical Engineering East ATTN: Dr. M. F. Zabielski University Park, Pennsylvania 16802
3	Gulf General Atomic ATTN: Dr. J. McGowan Dr. B. Turner Mr. J. Rutherford P. O. Box 608 San Diego, California 92112	1	The Rand Corporation ATTN: Dr. F. Gilmore 1700 Main Street Santa Monica, California 90406
2	General Electric Company Space Science Laboratory ATTN: Dr. M. Bortner Dr. R. Kummner P. O. Box 8555 Philadelphia, Pennsylvania 19101	2	Sandia Corporation ATTN: Dr. B. Long, Jr. Dr. G. Tisone P. O. Box 5800 Albuquerque, New Mexico 87115
3	Georgia Institute of Technology Physics Department ATTN: Dr. D. Martin Dr. E. McDaniel Dr. E. Thomas Atlanta, Georgia 30332	2	Southwest Center for Advanced Studies ATTN: Dr. C. Collins Dr. W. Hart P. O. Box 30365 Dallas, Texas 75230
		2	Stanford Research Institute ATTN: Dr. I. Poppoff Dr. F. Smith 333 Ravenswood Avenue Menlo Park, California 94025

DISTRIBUTION LIST

<u>No. of Copies</u>	<u>Organization</u>
1	University of Delaware Department of Physics ATTN: Dr. S. B. Woo Newark, Delaware 19711
2	University of Illinois Aeronomy Laboratory ATTN: Dr. C. Sechrist, Jr. Dr. S. Bowhill Urbana, Illinois 61801
4	University of Pittsburgh Cathedral of Learning ATTN: Prof. M. Biondi Prof. W. Fite Prof. F. Kaufman Dr. E. Zipf Pittsburgh, Pennsylvania 15213
2	University of Washington Department of Physics ATTN: Dr. R. Geballe Dr. K. Clark Seattle, Washington 98105

Aberdeen Proving Ground

Ch, Tech Lib
Marine Corps Ln Ofc
CDC Ln Ofc

UNCLASSIFIED

Security Classification

DOCUMENT CONTROL DATA - R & D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) U. S. Army Aberdeen Research and Development Center Ballistic Research Laboratories Aberdeen Proving Ground, Maryland 21005		2a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED	
3. REPORT TITLE NEGATIVE ION REACTIONS IN NO-H ₂ O MIXTURES		2b. GROUP	
4. DESCRIPTIVE NOTES (Type of report and Inclusive dates)			
5. AUTHOR(S) (First name, middle initial, last name) Lawrence J. Puckett and William C. Lineberger			
6. REPORT DATE December 1969		7a. TOTAL NO. OF PAGES 29	7b. NO. OF REFS 9
8a. CONTRACT OR GRANT NO.		9a. ORIGINATOR'S REPORT NUMBER(S) Report No. 1463	
b. PROJECT NO. RDT&E 5910.21.61160		9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
c.			
d.			
10. DISTRIBUTION STATEMENT This document has been approved for public release and sale; its distribution is unlimited.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY U. S. Army Materiel Command Washington, C. C.	
13. ABSTRACT A stationary afterglow system has been utilized to determine rate constants for thermal energy negative ion-molecule reactions in photoionized NO-H ₂ O mixtures. When the decay of the plasma is controlled by ambipolar diffusion of positive and negative ions quantitative determination of rate constants is shown to be feasible. The plasma transition from electron-positive ion ambipolar diffusive domination of the transport loss processes to domination by positive ion-negative ion ambipolar diffusion is explained by a model which includes the effects of negative ion trapping. Prominent negative ions in the afterglow include NO ₂ ⁻ , its hydrates, and clusters involving HNO ₂ . Reaction rate constants for the processes $\text{NO}_2^- + \text{H}_2\text{O} + \text{NO} \rightarrow \text{NO}_2^- \cdot \text{H}_2\text{O} + \text{NO}$ $\text{Cl}^- + \text{H}_2\text{O} + \text{NO} \rightarrow \text{Cl}^- \cdot \text{H}_2\text{O} + \text{NO}$ are found to be $1.3 \pm 0.3 \times 10^{-28} \text{ cm}^6/\text{sec}$ and $3.4 \pm 1.3 \times 10^{-29} \text{ cm}^6/\text{sec}$ at 293 K, respectively. Steady glows in NO-H ₂ O-O ₂ mixtures revealed that NO ₂ ⁻ and the impurity HCO ₃ ⁻ also formed multiple hydrates and clustered with HNO ₂ . These results indicate that the terminal negative ions in the D-region of the ionosphere will likely be hydrated.			

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

UNCLASSIFIED

Security Classification

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
Afterglows Nitric Oxide Hydrated Ions Hydronium Ions Chemical Reactions Reaction Rate Constants Clustered Ions Photoionization Negative Ions Ambipolar Diffusion Ionospheric Chemistry						

Security Classification